


⑦ Singular Value Decomposition ("SVD")

Given: data points " \vec{x}_i "




Least squares: Can find line

that minimizes $d_1^2 + d_2^2 + d_3^2 + d_4^2$

BUT: Only vertical distance

Sometimes that's the right thing - but
not always.



What if we'd like to find a line
to minimize

$$\|\vec{d}_1\|^2 + \|\vec{d}_2\|^2 + \|\vec{d}_3\|^2 + \|\vec{d}_4\|^2 ?$$

[Some answer as for least
squares line fitting problem? No.
see p.c.]

(Klein book: "trolley line problem")

Make problem easier (for now): Find line through origin that
minimizes ℓ_1 -norm distance
to (\vec{d}_i) .

Mathematically: Find direction \vec{v} (assume $\|\vec{v}\|_2 = 1$) s.t.

$$\sum_i \|\vec{d}_i\|_2^2 \quad (\text{where } \vec{d}_i \perp \vec{v}, \vec{x}_i - \vec{d}_i \in \text{span}\{\vec{v}\})$$

is minimized.

(Now drop vector arrows: $x_i = \vec{x}_i$, $d_i = \vec{d}_i$, $v = \vec{v}$)


[What is d_i ? $x_i^{\perp v}$]

$$x_i = x_i^{\perp v} + x_i^{\parallel v}$$

$$x_i = x_i^{\perp v} + (x_i, v)v$$

$$\xrightarrow{\text{Pyth}} \|x_i\|^2 = \|x_i^{\perp v}\|^2 + (x_i, v)^2 \|v\|^2$$

$\underbrace{}_{=1}$



$$\rightsquigarrow \sum_i \|d_i\|_2^2 = \sum_i \|x_i\|^2 - (x_i, v)^2$$

want to
minimize

$$= \|X\|_F^2 - \|Xv\|_2^2$$

\uparrow
Frobenius

\leftarrow Let $X = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$.

[How do we minimize that? Maximize $\|Xv\|_2^2$.]

[What's the maximal value of $\|Xv\|_2^2$? $\|X\|_2^2$]

[How can we find v ? Unclear (for now). Let's assume we can.]

Terminology

v : first right singular vector of $X \rightsquigarrow$ rename to v ,

$\sigma_1 := \|Xv\|_2$ first singular value of X

lec 16

review right / left singular vectors
relate to trolley line example

Q's:

review quiz

Next, assume we look for a vector $v_2 \perp v_1$, such that

$$\|Xv_2\|_2^2 \text{ is maximized.}$$

That's the second right singular vector of X .

And $\|Xv_2\|$ is the second singular value.

Next, assume we look for a vector $v_3 \perp v_1, v_2 \dots$

Then: $V^T = \begin{pmatrix} -v_1 & - \\ \vdots & \\ -v_n & - \end{pmatrix}$ if X is an $m \times n$ matrix. (assume that for now)

$\Sigma = \begin{pmatrix} \sigma_1 & & \\ & \sigma_2 & \\ & & \ddots & \sigma_n \end{pmatrix}$ where Σ is an $m \times n$ matrix in general.
↑ some shape as X
not a sum - the Greek letter Σ .

$$X = U \Sigma V^T$$

$$\rightsquigarrow X = U \Sigma V^T \quad \text{Derm: Finding the SVD}$$

Result: The singular value decomposition factors any $m \times n$ matrix
into:

$$X = U \Sigma V^T$$

where

- U is $m \times m$ orthogonal ← columns: "left singular vectors"
- Σ is diagonal, $m \times n$ and has positive entries
- V is $n \times n$ orthogonal ← columns: "right singular vectors"

Numerical example

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} 15 & 0 \\ 0 & 3 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -\frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & -\frac{\sqrt{3}}{2} \end{pmatrix}$$

$\underbrace{\qquad\qquad}_{\perp}$

Pseudo inverse

Define the pseudo inverse D^+ of an $m \times n$ diagonal matrix D as

$$D^+ = \begin{pmatrix} \frac{1}{\sigma_1} & & \\ & \frac{1}{\sigma_2} & \\ & & \ddots & \frac{1}{\sigma_{\min(m,n)}} \\ & 0 & & \end{pmatrix} \quad \text{for } D = \begin{pmatrix} \sigma_1 & & 0 \\ 0 & \ddots & 0 \\ 0 & & \sigma_n \end{pmatrix}$$

$$D^+ = \begin{pmatrix} \frac{1}{\sigma_1} & & 0 \\ & \frac{1}{\sigma_2} & \\ & & \ddots & \frac{1}{\sigma_{\min(m,n)}} \\ & 0 & & \end{pmatrix} \quad \text{for } D = \begin{pmatrix} \sigma_1 & 0 & \\ 0 & \ddots & 0 \\ 0 & & \sigma_n \end{pmatrix}$$

[What to do about zero (or near-zero) σ_i ? leave them as 0]

Then define the pseudo inverse A^+ of a general matrix A by its SVD.

If $A = U \Sigma V^T$, then define $A^+ = V \Sigma^+ U^T$.

[Pseudo inverse of example?]

Demo: Comparing the cost of LU, QR, SVD

lec 17

outline:

Q' 's: eigenvalues of $A =$ eigenvalues of A^T ?

SVD apps

SVD h-class quiz

quit review

eigval

practice exam submissions

review quiz

Applications of the SVD

① Least squares problems

Fact: $A^+ b$ solves the least squares problem $\min_x \|Ax - b\|_2$

This generalizes our QR-based method. $\rightarrow \text{HW}$

[How? By also allowing underdetermined systems. (Fewer eqns than unknowns, $A = \boxed{\text{fat}} \rightarrow \boxed{\text{short}}$)]

② Principal component analysis $\rightarrow \text{HW3}$

③ Computing $\|A\|_2$ $\|A\|_2 = \sigma_1$

④ Computing the 2-norm condition number

Assume A invertible.

Recall: $\kappa(A) = \|A\| \|A^{-1}\| = \sigma_1 / \sigma_n$

$$A = \begin{pmatrix} \|\cdot\|_2 \\ \vdots \\ \|\cdot\|_2 \end{pmatrix} \begin{pmatrix} \sigma_1 & & & \\ & \ddots & & \\ & & \ddots & \\ & & & \sigma_n \end{pmatrix} \begin{pmatrix} \overline{\Xi} \\ \Xi \\ \vdots \\ \Xi \end{pmatrix}$$

$$A^{-1} = \begin{pmatrix} \|\cdot\|_2 \\ \vdots \\ \|\cdot\|_2 \end{pmatrix} \begin{pmatrix} \frac{1}{\sigma_1} & & & \\ & \ddots & & \\ & & \frac{1}{\sigma_n} & \\ & & & \frac{1}{\sigma_n} \end{pmatrix} \begin{pmatrix} \overline{\Xi} \\ \Xi \\ \vdots \\ \Xi \end{pmatrix}$$

$\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n \geq 0$

Now, for any matrix (not just square/invertible), it turns out that


σ_1 / σ_n is the condition number.

(And that's also how it's computed. — Quite expensive!)

⑤ Low-rank approximation

$$A = \left(\begin{array}{c} \parallel u \parallel \\ \parallel v \parallel \end{array} \right) \left(\begin{array}{c} \sigma_1 \\ \sigma_2 \\ \vdots \\ \sigma_n \end{array} \right) \left(\begin{array}{c} \overline{v^T} \\ \overline{u^T} \end{array} \right)$$

$\left(\begin{array}{c} \sigma_1 u_1 \\ \sigma_2 u_2 \\ \vdots \\ \sigma_n u_n \end{array} \right)$

sum of outer products
 = sum of rank-1 matrices


$- \left(\begin{array}{c} 1 \\ u_1 u_2 \dots u_n \end{array} \right)$

$\sigma_1 u_1 v^T + \sigma_2 u_2 v^T + \dots + \sigma_n u_n v^T$

Idea: Could just use first few as an approximation.


In fact: $\sum_{i=1}^k \sigma_i u_i v^T$ is the closest rank-k matrix to A
 (measured in the Frobenius-norm.)

Demo: image compression

[What happens if $\text{rank } A < k$? $A = \sum_{i=1}^k \sigma_i u_i v^T$!]


[How does this relate to the trolley line problem? Closest rank-1!]

Original trolley line problem:



Line through the origin

More generally - how about:



Line not through origin?

Just need one point on that line:

Can then subtract that point from all data.

Fact: $\frac{1}{n} \sum_{i=1}^n x_i$ is such a point.

("centroid")

[Relationship to PCA?]

Lec 18

- exam 2

nullsp thru end of lecture
today


outline

hw3 due (ext: ask prof. DL)

hw 4 out Wed

Q:

Can define a variant of the SVD where U, V are not square.



"thin" SVD

[Why is this important?
 U consumes a large amount of
memory.